DOCUMENT RESUME

ED 406 442 TM 026 421

AUTHOR Rennie, Kimberly M.

TITLE Understanding the Sampling Distribution: Why We

Divide by n-1 To Estimate the Population Variance.

PUB DATE Jan 97

NOTE 20p.; Paper presented at the Annual Meeting of the

Southwest Educational Research Association (Austin,

TX, January 23-25, 1997).

PUB TYPE Reports - Descriptive (141) -- Speeches/Conference

Papers (150)

EDRS PRICE MF01/PC01 Plus Postage.

DESCRIPTORS Analysis of Variance; *Estimation (Mathematics);

*Sample Size; *Sampling; *Statistical Distributions;

Statistical Significance

IDENTIFIERS Population; Variance (Statistical)

ABSTRACT

This paper explains the underlying assumptions of the sampling distribution and its role in significance testing. To compute statistical significance, estimates of population parameters must be obtained so that only one sampling distribution is defined. A sampling distribution is the underlying distribution of a statistic. Sampling distributions are theoretical distributions that comprise an infinite number of sample statistics from an infinite number of randomly selected samples of a specified sample size. The influence that a large sample size has on statistical significance is demonstrated through some "what if" analyses. A "what if" analysis is simply an analysis of variance summary table in which the sample size is changed to see how statistical significance is affected. A large enough sample size invariably leads to statistical significance. Researchers with large sample sizes should look for other ways to interpret their results. One such way is effect size, which is a variance accounted for statistic that can tell how much of the variability in a dependent variable can be explained by the independent variables. (Contains 2 tables, 4 figures, and 12 references.) (SLD)



U.S. DEPARTMENT OF EDUCATION Office of Educational Research and Improvement EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

- This document has been reproduced as received from the person or organization originating it.
- Minor changes have been made to improve reproduction quality.
- Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.

PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL HAS BEEN GRANTED BY

KIMBERLY RENNIE

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

Understanding the Sampling Distribution: Why We Divide by n-1 to Estimate the Population Variance

Kimberly M. Rennie
Texas A&M University 77843-4225



Paper presented at the annual meeting of the Southwest Educational Research Association, Austin, January, 1997.

Often times, graduate students (especially those in the behavioral sciences) view statistics courses as classes in which they just have to get through. There is no desire to actually learn the material. Instead, students opt to memorize enough formulas to get a passing grade. As a result of this belief, when these graduate students take a statistics course, there is not any thinking involved. That is, they willingly accept what is being taught to them as the absolute and complete truth. Unfortunately, not all that is taught in courses or printed in Many dissertations (and research articles) books is true. contain methodological and design flaws. In fact, Thompson (1994) a paper about the seven common mistakes found dissertations. One mistake made by both graduate students and alike involves the interpretation of statistical faculty significance testing.

Significance Testing

The use of statistical significance testing in behavioral science research has been the subject of heated debate over the past two decades (Carver, 1978; Cohen, 1994; Greenwald, 1975; Thompson, 1993). More recently, the American Psychological Association (APA) has established a Task Force on Statistical Inference to consider banning the reporting of statistical significance testing in APA journals (Shea, 1996). Despite the efforts of APA and many notable researchers who argue against the improper use of statistical significance testing as the



determinant for declaring the results of a study important (Cohen, 1994; Thompson, 1989b), many researchers still rely solely on the use of statistical significance testing to claim that their findings are noteworthy (Kaminski & Good, 1996; Patel, Power, & Bhavnagri, 1996). Obviously, these researchers are not aware of the erroneous assertions that they are making. Thus, the first part of this paper will enlighten these researchers and others who are in danger of one day falling prey to the same fate by explaining, that statistical significance testing is driven in large part by sample size.

Sample size

Although, there are many reasons to argue against the use of statistical significance testing, the impact that sample size has on statistical significance testing seems to be the most salient way of demonstrating this point. "What if" analyses will be used to demonstrate how sample size directly impacts statistical significance testing (see Thompson, 1989a). A "what if" analysis is simply an ANOVA summary table in which the sample size is changed in order to see how statistical significance is affected by sample size, Tables 1 and 2 present these illustrations.

As (hopefully) all researchers know, if a sample is large enough obtaining statistically significant results is inevitable. Thompson (1996) noted that:

statistical significance testing primarily



'virtually any study can be made to show
[statistically] significant results if one
uses enough subjects' (Hays, 1981, p. 293)
As Nunnally (1960, p. 643) noted some 35 years
ago, 'If the null hypothesis is not rejected,
it is usually because the N is too small. If
enough data are gathered, the null hypothesis
will generally be rejected.' The implication is
that:

Statistical significance testing can involve a tautological logic in which tired researchers, having collected data from hundreds of subjects, then conduct a statistical test to evaluate whether there were a lot of subjects, which the researchers already know, because they collected the data and they know they're tired. This tautology has created considerable damage as regards the cumulation of knowledge. (Thompson, 1992, p. 436)

There is not an established method for determining the correct number of subjects that should be used in an experiment. Investigators can collect data from as few or as many subjects as they choose. Thus, conscientious researchers who collect data from a relatively large number of subjects will tend to obtain



5

statistically significant results regardless of the hypothesis that they are testing. This is demonstrated by the following example.

Assume that the sum-of-squares total is 100 and the data are analyzed with a one-way ANAOVA. An eccentric researcher out to find support for the hypothesis that people who eat apple pie have higher IQs than people who eat cherry pies. Obviously this hypothesis is pretty absurd-every self-respecting behavioral scientist knows that people who eat cherry pies are the ones with the higher IQs. However, this experimenter can obtain statistical significance for this hypothesis at a sample size of n=77 at p<.05, as noted in Table 2. Note that the effect size is only this scenario, this indicates that five percent. In preference can only account for five percent of the variance in IQ. An effect of this magnitude is not considered particularly large, according to Cohen's standards. Unfortunately, researcher who falsely believes that statistical significance testing measures how important results are will foolishly accept and attempt to publish these findings as noteworthy.

To further see the effects of sample size on statistical significance testing, different sample sizes were entered into a "what if" equation. In all of the examples the effect size was held constant. The results clearly show that as the sample size increases, \underline{F} calculated increases thereby making the probability



6

of statistically significant results more likely, as illustrated in Table 1.

Sampling distribution

In order to compute statistical significance, estimates of population parameters must first be obtained so that only one sampling distribution is defined (i.e., so that the sampling distribution is not statistically "indeterminate") (Thompson, 1996). Hence, the second part of this paper will explain the sampling distribution and the four properties of parameter estimates.

A sampling distribution is the underlying distribution of a statistic. Sampling distributions are theoretical distributions that are comprised of an infinite number of sample statistics taken from an infinite number of randomly selected samples of a specified sample size. For instance, if a random sample of size n=20 were taken from the population an infinite number of times, the combined means taken from all the samples would make up the sampling distribution of the mean. The ratio of the sample statistic (e.g., the mean of one sample of sample size n = 20) to the standard error of the statistic (i.e., the standard deviation statistic's sampling distribution) produces of statistics (e.g., t, F). These test statistics are then compared to the calculated values of the test statistics to determine if the results obtained are statistically significant.



For example, if an IQ test were given to a random sample of 100 graduate students and another to a random sample 100 of high school seniors, it is highly unlikely that the variance for the two sets of IQ scores would be the same. It is equally unlikely that either score would represent the actual population variance. Instead, these statistics would be estimates of the population variance. However, since the sample variance would tend to underestimate the actual population variance, a statistical correction (i.e., n-1) must be used in an attempt to correct for this bias. This bias and correction for this bias will be explained in more detail in the next section of the present paper.

Parameter Estimates

Parameter estimates have four properties: (a) unbiasedness, (b) consistency, (c) efficiency, and (d) sufficiency (Harnett, 1970). The properties of estimates of the population mean and estimates of the population variance will be utilized in order to explain these concepts.

Biasedness

Bias occurs when the difference between the parameter estimate and the population parameter is not equal to zero. A parameter estimate can accurately estimate, underestimate, or overestimate, the actual population parameters. In Figure 1, the parameter estimate (\underline{X}) perfectly estimates the actual population parameter (μ). This indicates that the parameter estimate is



equal to the actual population parameter and the estimate is unbiased (e.g., $\underline{X}=\mu$, where \underline{X} is the mean estimate and μ is the population mean). Figure 2 shows an underestimate of the population parameters. In this case, the parameter estimate is less than the population parameter (e.g., $SD^2 < \sigma^2$, where SD^2 is the sample variance and σ^2 is the population variance). When the parameter estimate is greater than the population parameter (e.g., y < Y, where y represents the parameter estimate and Y is the population parameter), this results in an overestimate of the parameter, as shown in Figure 3.

It is important to note that the mean estimate is always an unbiased estimate of the population mean and the variance estimate always underestimates the population variance. The following formula proves this fact for the mean (Harnett, 1970, p.159):

Define
$$\underline{X} = (1/n) (x_1 + x_2 + x_3 + ... + x_n)$$

$$E(\underline{X}) = E[(1/n) (x_1 + x_2 + x_3 + ... + x_n)]$$

$$= 1/nE[x_1 + x_2 + x_3 + ... + x_n]$$

$$= 1/n (E[x_1] + E[x_2] + E[x_3] + ... + x_n$$

$$= 1/n (\mu + \mu + \mu + ... + \mu)$$

$$= 1/n (n\mu)$$

$$\mu \underline{X} = \mu$$

$$\therefore EX = \mu$$



In random sample taken from a randomly distributed population, every person in the population has an equal chance of being selected. However, every score in the population does not have an equal chance of being selected. In a randomly distributed population, extreme scores have a lower probability of being selected, as illustrated in Figure 4. In this figure, it can be seen that the extreme scores have a 1 in 16 chance of being selected versus scores at the mean which have a 1 in four chance of being selected. Thus, extreme score will tend to underrepresented in the random sample. This results in the sample variance being lower than the variance in the population. In order to correct for this bias when calculating the variance, the SOS is divided by n-1 instead of n, which results in a larger result than when dividing b n-1.

Consistency

Consistency is the tendency of parameter estimates to become closer to the actual population parameter as the sample size increases. This occurs because it is expected that as sample size increases, the sample taken from the population becomes more representative of the population. Moreover, as sample size increases, the standard error of the statistic decreases (see Hinkle et al., 1994). Therefore, the sample statistics should become closer to the actual population values. The central limit theorem states that:

as sample size (n) increases, the sampling



distribution of the mean for simple random samples of n cases, taken from the population with a mean of μ and a finite variance equal to σ^2 , approximates a normal distribution.

(Hinkle et al., 1994, p. 150)

This is also true of the variance.

Efficiency

Efficiency has to do with the credibility of parameter estimates (e.g., how reliable is the estimate?). If two estimates are unbiased, the estimate which has the smaller variance in its' sampling distribution is more efficient (see Figures 5 & 6; Mittag, 1992).

Since the mean estimate is unbiased (i.e., the mean estimate is equal to he population mean), it will also be efficient. The variance, on the other hand, is never unbiased. As a result, the variance estimate is never 100% efficient. However, as the sample size increases, the variance estimate will become more efficient.

Sufficiency

Harnett, 1970 (p. 193) defined sufficiency as an estimator that "utilizes all of the information about the population parameter that is contained in the sample data." For example, the mode, median, and range represents estimates that are not sufficient. In both the sample and the population, the mode is the most common number in the distribution, the median is the number which divides the distribution into halves having an equal



number of persons or scores in the set of ordered scores, and the range is the highest number minus the lowest number in the distribution. In all of these cases, only one or two scores are used. Meanwhile, the mean, standard deviation, and the variance are all estimates that are sufficient. The following formulas demonstrate this:

sample population

 $mean = \Sigma x/n \qquad \mu = \Sigma x/N$

SD = (x-X)/n-1 $\sigma = (x-\mu)/N-1$

 $SD^2 = \sqrt{(x-X)/(n-1)}$ $\sigma^2 = \sqrt{(x-\mu)/(N-1)}$

Please note that in each of the preceding formulas every score in the distribution is utilized thereby fulfilling the requirements of being sufficient.

Conclusion

This paper explained the underlying assumptions behind the sampling distribution and its role in significance testing.

Moreover, the influence that a large sample size has on statistical significance was demonstrated through "what if" analyses. A large enough sample size invariably leads to statistical significance. Researchers with large sample sizes should look for other ways to interpret their results. One such way is effect size. Effect size is a variance accounted for statistic which can tell you how much of the variability in your dependent variable can be explained by your independent variable(s).



References

Cohen, J. (1994). The earth is round (\underline{p} <.05). American Psychologist, 45, 1304-1312.

Carver, R. (1978). The case against statistical significance testing. Harvard Educational Review, 48, 378-399.

Greenwald, A. (1975). Consequences of prejudices against the null hypothesis. Psychological Bulletin, 82, 1-20.

Harnett, D. L. (1970). <u>Introduction to statistical methods</u>
Reading, MA: Addison-Wesley Publishing Company.

Hinkle, D. E., Wiersma, W., & Jurs, S. G. (1994). Applied statistics for the behavioral sciences (3rd ed.). Boston, MA: Houghton Mifflin Company.

Kaminski, R. A., & Good, R. H., III. (1996). Toward a technology for assessing basic early literacy skills. School Psychology Review, 25(2), 215-227.

Patel, N., Power, T.G., & Bhavnagri, N. P. (1996).

Socialization values and practices of Indian immigrant parents:

Correlates of Modernity and acculturation. Child Development,

67(2), 302-313.

Shea, C. (1996). Psychologists debate accuracy of "significance test." Chronicle of Higher Education, 42(49), A12, A16.

Thompson, B. (1989a). Asking "what if" questions about significance tests. Evaluation in Counseling and Development, 22, 66-68.



Thompson, B. (1989b). Statistical significance, result importance, and result generalizability: Three noteworthy but somewhat different issues. Measurement and Evaluation in Counseling and Development, 22, 2-5.

Thompson, B. (1993). Theme issue: Statistical significance testing in contemporary practice. <u>Journal of Experimental</u> Education, 61(4).

Thompson, B. (1994, April). <u>Common methodology mistakes in dissertations</u>, <u>revisited</u>. Paper presented at the annual meeting of the American Educational Research Association, New Orleans. (ERIC Document Reproduction Service No. ED 368 771)



Table 1

Statistical Significance as Sample Size Increases

not significant-fail to reject the null

SOURCE	sos	DF	MS	F Calc	E.S.
residual	25	1	25	2.666667	0.25
error	75	8	9.375		
TOTAL	100	9	11.11111		

not significant-fail to reject the null

SOURCE	SOS	DF	MS	F Calc	E.S.
residual	25	1	25	4.333333	0.25
error	75	13 5.	769231		
TOTAL	100	14 7.	142857		

significant-reject the null

SOURCE	SOS	DF	MS	F Calc	E.S.
residual	25	1	25	6	0.25
error	75	18 4.	166667		
TOTAL	100	19 5.	263158		

significant-reject the null

SOURCE	SOS	DF	MS	F Calc	E.S.
residual	25	1	25	7.666667	0.25
error	75	23	3.26087		
TOTAL	100	24 4	1.166667		



Table 2

Statistical Significance Results for Cherry Pie Example significant-reject the null

SOURCE	sos	DF	MS	F Calc	E.S.
residual	5	1	5	4	0.05
error	95	76	1.25		
TOTAL	100	77 1.	298701		



Figure 1
Unbiased Estimate






Figure 2
Underbiased Estimate








Figure 3
Overbiased Estimate



y Y



Figure 4
Probability of Selecting Extreme Scores





TM 026421



U.S. DEPARTMENT OF EDUCATION

Office of Educational Research and Improvement (OERI) Educational Resources information Center (ERIC)



REPRODUCTION RELEASE

(Specific Document)

DOCUMENT IDENTIFICATION:

Title:	UNDERSTANDING THE SAMPLING DISTRIBUTION: WHY WE ESTIMATE THE POPULATION VARIANCE	DIVIDE BY n-1 TO
Autho	OTISI KIMBERLY M. RENNIE	
Corpo	orate Source:	Publication Date:
		1/97
11.	REPRODUCTION RELEASE: In order to disseminate as widely as possible timely and significant materials of announced in the monthly abstract journal of the ERIC system. Resources in Ex	of interest to the educational community, documents

in microfiche, reproduced paper copy, and electronic/optical media, and sold through the ERIC Document Reproduction Service (EDRS) or other ERIC vendors. Credit is given to the source of each document, and, if reproduction release is granted, one of the following notices is affixed to the document.

If permission is granted to reproduce the identified document, please CHECK ONE of the following options and sign the release

$\mathbf{X} \leftarrow$	Sample sticker to be affixed to document	Sample sticker to be affixed to document	
Check here	"PERMISSION TO REPRODUCE THIS MATERIAL HAS BEEN GRANTED BY	"PERMISSION TO REPRODUCE THIS MATERIAL IN OTHER THAN PAPER COPY HAS BEEN GRANTED BY	or here
(4"x 6" film). paper copy.	KIMBERLY M. RENNIE	<u>sample</u>	in other that
electronic. and optical media reproduction	TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)."	TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC):"	paper copy.

Level 1

Level 2

Sign Here, Please

Documents will be processed as indicated provided reproduction quality permits. If permission to reproduce is granted, but nertner box is checked, documents will be processed at Level 1.

"I hereby grant to the Educational Resources Information Center (ERIC) nonexclusive permission to reproduce this document as indicated above. Reproduction from the ERIC microfiche or electronic/optical media by persons other than ERIC employees and its system contractors requires permission from the copyright holder. Exception is made for non-profit reproduction by libraries and other service agencies to satisfy information needs of educators in response to discrete induiries."					
Signature: Kimberly lenne	Position: RESEARCH ASSOC				
Printed Name: KIMBERLY M. RENNIE	Organization: TEXAS A&M UNIVERSITY				
Address: TAMU DEPT EDUC PSYC	Telephone Number: (409) 845-1831				
COLLEGE STATION, TX 77843-4225	Date: 1/29/97				



III. DOCUMENT AVAILABILITY INFORMATION (FROM NON-ERIC SOURCE):

If permission to reproduce is not granted to ERIC. or, if you wish ERIC to cite the availability of this document from another source, please provide the following information regarding the availability of the document. (ERIC will not announce a document unless it is publicly available, and a dependable source can be specified. Contributors should also be aware that ERIC selection criteria are significantly more stringent for documents which cannot be made available through EDRS).

Publisher/Distributor:	
Address:	
	<u> </u>
Price Per Copy:	Quantity Price:
. REFERRAL OF ERIC TO	COPYRIGHT/REPRODUCTION RIGHTS HOLDER:
If the right to grant reproduction rel name and address:	riease is neid by someone other than the addressee. Diease provide the appropriate
Name and address of current copyright/reproc	duction rights halder:
Name:	
Address:	
•	
. WHERE TO SEND THIS	S =====
WHERE TO SEND THIS	5 FORM:
Send this form to the following ERIC Clearing	gnouse:

If you are making an unsolicited contribution to ERIC, you may return this form (and the document being contributed) to:

ERIC Facility 1301 Piccard Orive, Suite 300 Rockville, Maryland 20850-4305 Telephone: (301) 258-5500

